



## Total Participants

### Distribution by gender

Women	25	78%
Men	7	22%
<b>Total</b>	<b>32</b>	

(\*) Including Health Professionals, volunteers, and children

### Distribution by ages

<18	0	0%
18-28	12	39%
29-35	4	13%
36-45	4	13%
46-54	5	16%
>=55	6	19%
<b>Total</b>	<b>31</b>	

### Total Cholesterol

Optimal	<200	23	85%
Borderline	200-239	4	15%
At-risk	>=240	0	0%

### Non-fasting Glucose

# people

Low	< 65	0	0%
Optimal	65-139	22	100%
Borderline	140-199	0	0%
At-risk	>=200	0	0%

### Glucose - Fasting

Low	<65	0	0%
Optimal	65-100	2	40%
Borderline	101-124	2	40%
At-risk	>=125	1	20%

### Waist circumference

At risk men	>40	3	13%
At risk women	>35	9	39%
No at risk men	<=40	2	9%
No at risk wome	<=35	9	39%

### Blood Pressure

# people

Normal	<120	<80	0	0%
Borderline	120-139	80-89	16	76%
At-risk 1	140-159	90-99	4	19%
At-risk 2	>=160	>=100	1	5%

## LIPID PANEL

### 27 LDL - Bad Cholesterol

Optimal	<130	3	60%	5
Borderline	130-159	1	20%	
At-risk	>=160	1	20%	

22

### HDL - Good Cholesterol

Optimal	>=60	2	40%	5
Borderline women	41-59	2	40%	
At-risk women	<= 40	0	0%	
Borderline men	51-59	0	0%	
At-risk men	<= 50	1	20%	

5

### Triglycerides

Optimal	<150	4	80%	5
Borderline	150-199	1	20%	
At-risk	>=200	0	0%	

23

### BMI

# people

Underweight	<18.5	1	4%	23
Normal	18.5-24.9	6	26%	
Overweight	25 - 29.9	6	26%	
Obese	30- 39.9	7	30%	
Morbidly obese	>=40	3	13%	

### Explanation Results

# people

27

**TOTAL NUMBER OF SCREENINGS/SERVICES GIVEN**

**163**

