



Total Participants

Distribution by gender

Women	31	74%
Men	11	26%
Total	42	

(*) Including Health Professionals, volunteers, and children

Distribution by ages

<18	2	5%
18-28	17	40%
29-35	9	21%
36-45	5	12%
46-54	6	14%
>=55	3	7%

42

Total Cholesterol

Optimal	<200	22	81%
Borderline	200-239	4	15%
At-risk	>=240	1	4%

Glucose - Fasting

Low	<65	0	0%
Optimal	65-100	18	67%
Borderline	101-124	6	22%
At-risk	>=125	3	11%

BMI

Underweight	<18.5	0	0%
Normal	18.5-24.9	10	40%
Overweight	25 - 29.9	7	28%
Obese	30- 39.9	7	28%
Morbidly obese	>=40	1	4%

Blood Pressure

Normal	<120	<80	14	61%
Borderline	120-139	80-89	7	30%
At-risk 1	140-159	90-99	2	9%
At-risk 2	>=160	>=100	0	0%

Depression Screening

Less than 3 depression factors	9	90%
3 or more than 3 depression factors	1	10%

Waist circumference

At risk men	>40	0%	
At risk women	>35	0%	
Not at risk men	<=40	2	67%
Not at risk women	<=35	1	33%

Clinical Breast Exam

Normal	4	67%
Abnormal	2	33%

27

LIPID PANEL

LDL - Bad Cholesterol

Optimal	<130	24	89%
Borderline	130-159	2	7%
At-risk	>=160	1	4%

27

HDL - Good Cholesterol

Optimal	>=60	4	15%
Borderline women	41-59	8	30%
At-risk women	<= 40	7	26%
Borderline men	51-59	1	4%
At-risk men	<= 50	7	26%

25

Triglycerides

Optimal	<150	20	74%
Borderline	150-199	3	11%
At-risk	>=200	4	15%

27

BMI

Underweight	<18.5	0	0%
Normal	18.5-24.9	10	40%
Overweight	25 - 29.9	7	28%
Obese	30- 39.9	7	28%
Morbidly obese	>=40	1	4%

23

25

Explanation Results

people

27

Hemoglobin

27

Optimal women	11.7-15.5	15	56%
Low women	<11.7	4	15%
Optimal men	13.2-17.1	6	22%
Low men	<13.2	2	7%

6

TOTAL NUMBER OF SCREENINGS GIVEN

281