



Total Participants

Distribution by gender

| | | |
|--------------|-----------|-----|
| Women | 28 | 42% |
| Men | 39 | 58% |
| Total | 67 | |

(*) Including Health Professionals, volunteers, and children

Distribution by ages

| | | |
|-------|----|-----|
| <18 | 6 | 9% |
| 18-28 | 17 | 25% |
| 29-35 | 8 | 12% |
| 36-45 | 12 | 18% |
| 46-54 | 12 | 18% |
| >=55 | 12 | 18% |

67

| Total Cholesterol | | | |
|-------------------|---------|----|-----|
| Optimal | <200 | 42 | 81% |
| Borderline | 200-239 | 5 | 10% |
| At-risk | >=240 | 5 | 10% |

| Non-fasting Glucose | | | |
|---------------------|---------|----|-----|
| Low | < 65 | 0 | 0% |
| Optimal | 65-139 | 47 | 87% |
| Borderline | 140-199 | 5 | 9% |
| At-risk | >=200 | 2 | 4% |

| Hemoglobin | | | |
|---------------|-----------|----|-----|
| Optimal women | 11.7-15.5 | 15 | 29% |
| Low women | <11.7 | 5 | 10% |
| Optimal men | 13.2-17.1 | 31 | 60% |
| Low men | <13.2 | 1 | 2% |

52

54

52

BMI

| | | | |
|----------------|-----------|----|-----|
| Underweight | <18.5 | 1 | 2% |
| Normal | 18.5-24.9 | 17 | 31% |
| Overweight | 25 - 29.9 | 21 | 38% |
| Obese | 30- 39.9 | 13 | 24% |
| Morbidly obese | >=40 | 3 | 5% |

55

Blood Pressure

| | | | | |
|------------|---------|-------|----|-----|
| Normal | <120 | <80 | 18 | 37% |
| Borderline | 120-139 | 80-89 | 19 | 39% |
| At-risk 1 | 140-159 | 90-99 | 11 | 22% |
| At-risk 2 | >=160 | >=100 | 1 | 2% |

49

Depression Screening

| | | |
|-------------------------------------|----|-----|
| Less than 3 depression factors | 9 | 43% |
| 3 or More than 3 depression factors | 12 | 57% |

21

Explanation Results

people 58

TOTAL NUMBER OF SCREENINGS GIVEN

341