



### Total Participants

#### Distribution by gender

Women	160	62%
Men	97	38%
<b>Total</b>	<b>257</b>	

(\*) Including volunteers, and children

#### Distribution by ages

<18	12	5%
18-28	32	12%
29-35	53	21%
36-45	90	35%
46-54	40	16%
>=55	31	12%
	<b>258</b>	

<b>Total Cholesterol</b>				<b>207</b>
Optimal	<200	121	58%	
Borderline	200-239	53	26%	
At-risk	>=240	33	16%	

<b>Non-fasting Glucose</b>				<b>210</b>
Low	< 65	0	0%	
Optimal	65-139	177	84%	
Borderline	140-199	20	10%	
At-risk	>=200	13	6%	

<b>HDL - Good Cholesterol</b>				<b>156</b>
Optimal	>=60	21	13%	
Borderline men	41-59	28	18%	
At-risk men	<= 40	29	19%	
Borderline women	51-59	25	16%	
At-risk women	<= 50	53	34%	

<b>Albumin/Creatinin (A:C)</b>				<b>133</b>
Optimal	<30	122	92%	
Borderline	30 - 300	10	8%	
At-risk	>300	1	1%	

<b>Blood Pressure</b>					<b>202</b>
Normal	<120	<80	100	50%	
Borderline	120-139	80-89	78	39%	
At-risk 1	140-159	90-99	19	9%	
At-risk 2	>=160	>=100	5	2%	

<b>BMI</b>				<b>198</b>
Underweight	<18.5	0	0%	
Normal	18.5-24.9	24	12%	
Overweight	25 - 29.9	98	49%	
Obese	30- 39.9	73	37%	

<b>Waist circumference</b>					<b>193</b>
Optimal men	<=37	37	19%		
Optimal women	<=31.5	12	6%		
Borderline men	>37	23	12%		
Borderline women	>31.5	18	9%		
At risk men	>=40	21	11%		
At risk women	>=34.5	82	42%		

<b>Hemoglobin</b>				<b>180</b>
Optimal women	12-15.5	65	36%	
Low women	<12	38	21%	
Optimal men	13.5-17.5	59	33%	
Low men	<13.5	18	10%	

<b>Explanation Results</b>					<b>207</b>
# people					
<b>Pulse</b>					<b>182</b>
Normal	60-100	175	96%		
At-risk	<60	7	4%		

<b>Oxygen Saturation</b>				<b>182</b>
Normal	95-100	173	95%	
Borderline	90-94	9	5%	
At-risk	< 90	0	0%	

<b>A1C</b>				<b>24</b>
Optimal	<= 7%	15	63%	
At-risk	> 7%	9	38%	

<b>TOTAL NUMBER OF SCREENINGS GIVEN</b>		<b>2382</b>
---	--	-------------